

A1  
6. (Amended) The assay according to claim 1 wherein said assay is used to identify compounds that disrupt normal membrane receptor interactions, or can in themselves disrupt such interactions.

7. (Amended) The assay according to claim 1 for detecting a compound which serves as an inverse agonist, antagonist or agonist of the membrane receptor.

---

A2  
9. (Amended) The assay according to claim 1 wherein said membrane receptor is a growth factor receptor, cytokine receptor, ion channel, integrin, or G-protein receptor.

---

13. (Amended) The assay according to claim 9 wherein said G-protein coupled receptor is a dopamine receptor, a muscarinic cholinergic receptor, an  $\alpha$ -adrenergic receptor, a  $\beta$ -adrenergic receptor, an opiate receptor, a cannabinoid receptor, a serotonin receptor or a protease activated receptor.

A3  
14. (Amended) The assay according to claim 1 wherein the receptor/reporter fusion protein is expressed from nucleic acid construct comprising a gene encoding said reporter protein which is fused in-frame to the 5' or 3' end of a gene encoding said membrane receptor.

15. (Amended) The assay according to claim 1 wherein the functionality of said membrane receptor/reporter fusion protein is substantially unaffected by fusion of the reporter protein to the receptor.

16. (Amended) The assay according to claim 15 wherein said reporter protein is Green Fluorescent Protein (GFP), or active variant thereof.

---

A4  
18. (Amended) The assay according to claim 15 wherein said reporter protein is *Renilla reniformis* (sea pansy) luciferase protein, secreted placental alkaline phosphatase

A 4  
conce (SEAP),  $\beta$ -lactamase, galactosidase, firefly (*Photinus pyralis*) luciferase, blue fluorescent protein, yellow fluorescent protein, and cyan fluorescent protein.

20. (Amended) The assay according to claim 1 wherein said reporter protein is used to localize and/or quantify the membrane receptor.

21. (Amended) An assay according to claim 20 wherein any change of said membrane receptor/reporter fusion protein is detected as a change in cellular localisation of the receptor/reporter fusion protein, or semi-quantitatively by the synthesis or degradation of said receptor/reporter fusion protein.

22. (Amended) An assay according to claim 1 wherein said detection of any change of said membrane receptor/reporter fusion protein is carried out with cells placed on the surface of a microscope slide.

A 5  
23. (Amended) The assay according to claim 1 wherein said detection of any change of said membrane receptor/reporter fusion protein is carried out on cells placed in a well of a microtitre plate.

A 6  
29. (Amended) The membrane receptor/reporter fusion protein according to claim 28 wherein the reporter protein is GFP or luciferase.

Please add the following new claim.

A 7  
32. (Amended) The membrane receptor/reporter fusion protein according to claim 27 wherein the reporter protein is GFP or luciferase.